

UB-TH-0193
hep-th/9307175
Revised: March 1, 1994

Multi-Soliton Solutions of Affine Toda Models

Z. Zhu and D. G. Caldi

Department of Physics
State University of New York at Buffalo
Buffalo, N.Y. 14260-1500

ABSTRACT

Hirota's method is used to construct multi-soliton and plane-wave solutions for affine Toda field theories with imaginary coupling.

1 Introduction

There has been a revived interest in Toda field theories [1-12] recently, chiefly because they represent a large class of integrable two-dimensional models which include not only conformal field theories (CFT's) but also massive deformations away from conformality. Toda field theories (TFT's) have a Lagrangian realization (and hence provide one for the CFT's which can be written as TFT's) and can be classified according to their underlying Kac-Moody algebra g . If g is a finite-dimensional, semi-simple Lie algebra, the theory is a CFT. If g is an affine Kac-Moody loop algebra, the resulting affine Toda field theory is massive, but still integrable. Indeed, these theories provide explicit realizations of the phenomenon investigated by Zamolodchikov [13] concerning integrable massive deformations of CFT's. Finally, it has been observed [5, 7] that if one has the full affine Kac-Moody algebra (i.e., with central extension and, in fact, two more fields compared with the usual TFT), then the theory is conformal again and is called a conformal affine Toda field theory. The affine TFT is then considered a "gauge-fixed" version of the conformal affine theory.

In this paper we are concerned with finding exact solutions to classical affine TFT's. To this end we use the method introduced by Hirota [14] and applied to TFT's in the stimulating work of Hollowood [1]. If the coupling in the theory is real, then the Toda fields can be consistently taken as real fields. The theories are unitary and the S-matrices have been solved [15]. Since the interaction potential has a unique minimum, there are no solitons. But these real affine TFT's are not those related to unitary CFT's. Curiously enough, it turns out that to have the relationship to CFT's mentioned above, one must take the coupling constant to be purely imaginary [1]. But then one has the additional feature that the Toda potential is periodic, i.e., invariant under translations on the weight lattice of g . Hence, the vacua are infinitely degenerate and there is a rich spectrum of topological solitons. But the

Lagrangian is complex and the Hamiltonian is non-Hermitian; so the theory would appear to be non-unitary. Yet the solitons have real masses, and it has been suggested that for some region of the imaginary coupling, there may exist a truncation which would render the theories unitary [2].

Hirota's method has been successfully applied to the affine Toda theories with purely imaginary coupling. Single-soliton solutions were obtained in [1, 6, 7] and multi-soliton solutions for $A_r^{(1)}$ theories in [1]. In this paper, we will construct more multi-soliton solutions to the complex affine Toda theories.

In §2, we will review briefly the affine Toda theories and Hirota's method. Since the solutions for non-simply-laced Toda models can be derived from the simply-laced ones, we will mostly work on the simply-laced cases with some comments on non-simply-laced cases. In §3, we construct multi-soliton solutions for the $\hat{A}_r$ loop algebras, in §4, we deal with $\hat{so}(8)$, and in §5, $\hat{D}_{2r}$, We summarize our results and conclusions in §6.

2 Affine Toda Models and Hirota's Method

The Lagrangian density of affine Toda field theory can be written in the form

$$\mathcal{L} = \frac{1}{2}(\partial_\mu \varphi)(\partial^\mu \varphi) - \frac{2}{\psi^2} \frac{m^2}{\tilde{\beta}^2} \sum_{j=0}^r n_j (e^{\tilde{\beta} \alpha_j \cdot \varphi} - 1). \quad (2.1)$$

The field $\varphi(x, t)$ is an r -dimensional vector, r being the rank of some finite dimensional semi-simple Lie algebra g . The α_j 's, for $j = 1, \dots, n$ are the simple roots of g ; ψ is the highest root: $\psi = \sum_{j=1}^r n_j \alpha_j$, where the n_j 's are positive integers, and $n_0 = 1$. $\alpha_0 = -\psi$ is the extended root of $\hat{g}$, the affine extension of g . $\tilde{\beta}$ is the coupling constant, m is the mass parameter. Note, in general repeated indices are not summed over in this paper.

The equations of motion are

$$\partial^2 \varphi = -\frac{2}{\psi^2} \frac{m^2}{\tilde{\beta}^2} \sum_{j=0}^r n_j \alpha_j e^{\tilde{\beta} \alpha_j \cdot \varphi}. \quad (2.2)$$

Writing φ out in component form, $\varphi = \sum_{j=1}^r (2\alpha_j/\alpha_j^2) \varphi^j$, we have

$$\partial^2 \varphi^j = -\frac{m^2}{\tilde{\beta}} l_j \left(\exp(\tilde{\beta} \sum_{k=0}^r K_{jk} \varphi^k) - \exp(\tilde{\beta} \sum_{k=0}^r K_{0k} \varphi^k) \right), \quad (2.3)$$

where $K_{jk} = 2\alpha_j \cdot \alpha_k / \alpha_k^2$ is the the Cartan matrix of $\hat{g}$, and l_j 's are integers related to n_j 's by $l_j = (\alpha_j^2/\psi^2) n_j$, for $j = 0, 1, \dots, r$. As noted in the introduction, we are interested in the case where the coupling constant is purely imaginary,

$$\tilde{\beta} = i\beta.$$

We now use Hirota's method to solve the equations of motion. One introduces a nonlinear transformation

$$\varphi = -\frac{1}{i\beta} \sum_{j=0}^r \frac{2\alpha_j}{\alpha_j^2} \ln \tau_j, \quad \text{or} \quad \varphi^j = -\frac{1}{i\beta} (\ln \tau_j - l_j \ln \tau_0), \quad (2.4)$$

where the new variables are the tau-functions for the theories. The equations of motion (2.2) then decouple to a collection of equations in Hirota's form,

$$\frac{1}{2m^2} (D_t^2 - D_x^2) \tau_j \cdot \tau_j = l_j \left(\prod_{k=0}^r \tau_k^{-K_{jk}} - 1 \right) \tau_j^2. \quad (2.5)$$

Here D_x and D_t are Hirota derivatives, defined by

$$D_x^m D_t^n f \cdot g = \left(\frac{\partial}{\partial x} - \frac{\partial}{\partial x'} \right)^m \left(\frac{\partial}{\partial t} - \frac{\partial}{\partial t'} \right)^n f(x, t) g(x', t') \Big|_{\substack{x=x' \\ t=t'}}.$$

We expand the τ -functions in a formal power series in an arbitrary parameter ϵ as

$$\tau_j = \sum_{n=0} \epsilon^n \tau_j^{(n)} = 1 + \epsilon \tau_j^{(1)} + \epsilon^2 \tau_j^{(2)} + \dots. \quad (2.6)$$

Using the bilinearity of Hirota's operator, we can rewrite (2.5) as

$$\frac{1}{2m^2} \sum_n \sum_{a+b=n} \epsilon^n (D_t^2 - D_x^2) \tau_j^{(a)} \cdot \tau_j^{(b)} = l_j \left(\prod_{k=0}^r \tau_k^{-K_{jk}} - 1 \right) \tau_j^2. \quad (2.7)$$

Expanding the r.h.s. also as power series in ϵ , and equating corresponding coefficients, the resulting equation can be written in the form:

$$\sum_k L_{jk} \tau_k^{(n)} + \frac{1}{m^2} (\partial_t^2 - \partial_x^2) \tau_j^{(n)} = U_j^{(n)}, \quad (2.8)$$

where $L_{jk} = l_j K_{jk}$, and $U_j^{(n)}$ collects the rest of the n^{th} order terms which depend on $\tau^{(1)}, \dots, \tau^{(n-1)}$. The explicit form of $U_j^{(n)}$ depends on the Cartan matrix K . Now the nonlinear equations of motion are simplified to a set of linear partial differential equations, which can be solved iteratively. Furthermore, the first order equation is homogeneous, $U_j^{(1)} = 0$.

For a single soliton traveling at velocity v , τ_j should depend on x and t only through the combination $z = x - vt$, $\tau_j(x, t) = \tau_j(x - vt) = \tau_j(z)$. Here equation (2.8) simplifies to a set of ordinary differential equations:

$$\sum_k L_{jk} \tau_k^{(n)}(z) - \frac{1 - v^2}{m^2} \frac{d^2}{dz^2} \tau_j^{(n)}(z) = U_j^{(n)}, \quad (2.9)$$

with the first order equation given by:

$$(1 - v^2) \frac{d^2}{dz^2} \tau_j^{(1)}(z) = m^2 \sum_k L_{jk} \tau_k^{(1)}(z). \quad (2.10)$$

Equation (2.10) can be solved by diagonalizing the matrix L_{jk} . There are two linearly independent solutions for any given eigenvector Λ (with corresponding eigenvalue λ). For the null eigenvector $\Lambda_0 = (l_0, l_1, \dots, l_r)$, with eigenvalue $\lambda_0 = 0$, the two linearly independent solutions are:

$$\tau_j^{(1)}(z) = (c_1 + c_2 z) l_j, \quad (2.11)$$

which correspond to vacuum solutions. For Λ with non-zero eigenvalue λ , the two linearly independent solutions are:

$$\tau_j^{(1)}(z) = \Lambda_j e^{\gamma z + \xi}, \quad (2.12)$$

where ξ is an integration constant, and γ satisfies $(1 - v^2)\gamma^2 = m^2\lambda$. Since γ can be either positive or negative, (2.11) and (2.12) give all the linearly independent solutions to (2.10).

General solutions to (2.10) are superposition of the solutions given in (2.12), solution to (2.9) can also be superposed with a solution to (2.10), but they will not have properties of a single soliton, and will be dealt with later as multi-soliton solutions. Here we will restrict our solutions to the single-soliton form:

$$\tau_j^{(n)} = \delta_j^{(n)} e^{n\Gamma}, \quad (2.13)$$

where $\Gamma = \gamma(x - vt) + \xi$, and $\delta_j^{(n)}$ is independent of x and t . Now (2.9) can be simplified to a set of linear algebraic equations:

$$\sum_k (L_{jk} - n^2 \lambda \delta_{jk}) \delta_k^{(n)} = V_j^{(n)}, \quad (2.14)$$

where $V_j^{(n)}$ depends only on $\delta^{(1)}, \dots, \delta^{(n-1)}$. In particular, $V_j^{(1)} = 0$, so $\delta^{(1)}$ is an eigenvector of L with eigenvalue λ . $\delta^{(2)}, \delta^{(3)}, \dots$, can be solved from (2.14) uniquely, except for $su(6p)$ and $sp(6p)$, where additional degeneracy exists. In [7] it is shown that $\delta_j^{(n)}$ vanishes if $n > \kappa l_j$ for some integer κ , which is less than or equal to the degeneracy of the eigenvalue λ , except for the special cases of $su(6p)$ and $sp(6p)$ noted above.

Since the equation (2.8) for $\tau_j^{(1)}$ is linear and homogeneous, one can superpose the single-soliton solutions to get general N -soliton solutions, at least up to the first order:

$$\tau_j^{(1)} = \epsilon_1 \delta_{1,j}^{(1)} e^{\Gamma_1} + \dots + \epsilon_N \delta_{N,j}^{(1)} e^{\Gamma_N}, \quad (2.15)$$

where ϵ_a for $a = 1, \dots, N$ are some arbitrary expansion parameters, $\delta_{a,j}^{(1)}$ is the first order coefficient of the a^{th} soliton and is an eigenvector of the L matrix, as discussed above. For a general N -soliton solution, we use the following expansion for τ_j :

$$\tau_j = \sum_{\{m_a\}} \epsilon_1^{m_1} \dots \epsilon_N^{m_N} \delta_j(m_1, m_2, \dots, m_N) \exp(m_1 \Gamma_1 + m_2 \Gamma_2 + \dots + m_N \Gamma_N), \quad (2.16)$$

where $\delta_j(0, 0, \dots, 0) = 1$, and from (2.15)

$$\delta_j(0, \dots, 0, m_a = 1, 0, \dots, 0) = \delta_{a,j}^{(1)}. \quad (2.17)$$

From now on, we will let $\epsilon_a = 1$.

Let us define

$$\begin{aligned} f(m_1, m_2, \dots, m_N) &\equiv \frac{1}{m^2} \left((m_1 \gamma_1 + \dots + m_N \gamma_N)^2 - (m_1 \gamma_1 v_1 + \dots + m_N \gamma_N v_N)^2 \right) \\ &= \sum_a m_a^2 \lambda_{s_a} + 2 \sum_{a \neq b} m_a m_b \sqrt{\lambda_{s_a} \lambda_{s_b}} \cosh \theta_{ab}, \end{aligned} \quad (2.18)$$

where θ_{ab} is the rapidity difference of a^{th} and b^{th} soliton, and $\lambda^{(a)}$ is the eigenvalue for the a^{th} soliton, we have assumed that all the γ 's are positive just for definiteness. Using the the following property of Hirota's derivative operators:

$$(D_t^2 - D_x^2) e^{\alpha_1 x + \beta_1 t} \cdot e^{\alpha_2 x + \beta_2 t} = \left((\beta_1 - \beta_2)^2 - (\alpha_1 - \alpha_2)^2 \right) e^{(\alpha_1 + \alpha_2)x + (\beta_1 + \beta_2)t}, \quad (2.19)$$

we can write:

$$\begin{aligned} \frac{1}{2m^2} (D_t^2 - D_x^2) \tau_j \cdot \tau_j + l_j \tau_j^2 = \\ - \sum_{k_a} \left((f(k_1, \dots, k_N) - 2l_j) \delta_j(k_1, \dots, k_N) + A_j(k_1, \dots, k_N) \right) e^{k_1 \Gamma_1 + \dots + k_N \Gamma_N}, \end{aligned} \quad (2.20)$$

where

$$\begin{aligned} A_j(k_1, \dots, k_N) = \sum_{0 \prec (m_1, \dots, m_N) \prec (k_1, \dots, k_N)} \left(\frac{1}{2} f(k_1 - 2m_1, \dots, k_N - 2m_N) - l_j \right) \cdot \\ \delta_j(m_1, \dots, m_N) \delta_j(k_1 - m_1, \dots, k_N - m_N). \end{aligned}$$

The partial ordering $(m_1, \dots, m_N) \prec (k_1, \dots, k_N)$ is defined as $m_a \leq k_a$, for all a 's, and $\sum m_a < \sum k_a$. Also we can write:

$$\prod_{k \neq j} \tau_k^{-K_{jk}} = \sum_{k_1, \dots, k_N} \left(- \sum_{k \neq j} K_{jk} \delta_k(k_1, \dots, k_N) + B_j(k_1, \dots, k_N) \right) e^{k_1 \Gamma_1 + \dots + k_N \Gamma_N}, \quad (2.21)$$

where $B_j(k_1, \dots, k_N)$ is a function of the $\delta_j(m_1, \dots, m_N)$'s for $(m_1, \dots, m_N) \prec (k_1, \dots, k_N)$.

The explicit form of $B_j(k_1, \dots, k_N)$ will depend on the Cartan matrix K . Using (2.20) and (2.21), the nonlinear partial differential equation (2.5) can now be simplified to a set of linear algebraic equations:

$$\sum_k \left(L_{jk} - f(k_1, \dots, k_N) \delta_{jk} \right) \delta_k(k_1, \dots, k_N) = V_j(k_1, \dots, k_N), \quad (2.22)$$

where $V_j(k_1, \dots, k_N) = A_j(k_1, \dots, k_N) + l_j B_j(k_1, \dots, k_N)$. Since the r.h.s. of (2.22) only depends on the preceding δ_j 's, as discussed after (2.21), the above equation can be solved iteratively.

First we decompose the right hand side of (2.22) into linear combinations of the eigenvectors,

$$V_j(k_1, \dots, k_N) = \sum_s c_s(k_1, \dots, k_N) \Lambda_{s,j}, \quad (2.23)$$

where Λ_s , for $s = 0, 1, \dots, r$, is the complete set of eigenvectors of the matrix L . Since L is not necessarily symmetric, the eigenvectors are not necessarily orthogonal to each other with respect to the usual inner product. But for simply-laced loop algebras, eigenvectors of the matrix L corresponding to distinct eigenvalues are orthogonal to each other with respect to the diagonal metric $(1/l_0, 1/l_1, \dots, 1/l_N)$. This observation will make the decomposition (2.23) easier. Now from equations (2.22) and (2.23) one can see that

$$\delta_j(k_1, \dots, k_N) = \sum_s \frac{c_s(k_1, \dots, k_N) \Lambda_{s,j}}{\lambda_s - f(k_1, \dots, k_N)}. \quad (2.24)$$

Because of the iterative nature of (2.22), and from the initial choice (2.17), one can show that

$$\begin{aligned} \delta_j(0, \dots, 0, m_a, 0, \dots, 0) &= \delta_{a,j}(m_a), \\ \delta_j(0, \dots, 0, m_a, 0, \dots, m_b, \dots, 0) &= \delta_{ab,j}(m_a, m_b), \\ &\vdots \end{aligned} \quad (2.25)$$

where the $\delta_{ab,j}(m_a, m_b)$ is a coefficient from the pair scattering solution of the a^{th} and b^{th} soliton, etc. This means that the coefficients of $(N-1)$ -soliton solutions can be used in the N -soliton solutions, this property makes solving general N -soliton solutions tractable.

We conclude this section with a number of observations and remarks, the first of which concerns integrability. Due to it, the perturbative expansion terminates at finite order. By

counting the powers in ϵ , it is shown in [7] that, for a single-soliton solution, the highest nonvanishing order in τ_j is κl_j for some constant integer κ . Assuming that the expansion (2.16) gives exact solutions, then by repeating the power counting for each ϵ_a , one can show that:

$$\delta_j(m_1, \dots, m_N) = 0, \quad \text{if } (m_1, \dots, m_N) \not\leq (\kappa'_1 l_j, \dots, \kappa'_N l_j). \quad (2.26)$$

For all the double-soliton solutions we obtained, we can explicitly verify that $\kappa'_a = \kappa_a$ for $a = 1, 2$, where κ_a is the κ value for the a^{th} soliton. For a general N -soliton solution it is also true that $\kappa'_a = \kappa_a$ for $a = 1, 2, \dots, N$, since otherwise the resulting expression would not reduce to N single-soliton solutions in the asymptotic limit. Let $l_{\max} = \max(l_1, \dots, l_N)$, from the above discussion and (2.22) we get:

$$V_j(m_1, \dots, m_N) = 0, \quad \text{if } (m_1, \dots, m_N) \not\leq l_{\max} (\kappa_1, \dots, \kappa_N). \quad (2.27)$$

We consider a solution found if all nonvanishing δ_j 's are found.

If the Dynkin diagram of the loop algebra $\hat{g}_1$ has non-trivial automorphisms, then $\hat{g}_1$ can be twisted to form a new loop algebra $\hat{g}_2$. The equations of motion of the Toda theory associated to $\hat{g}_1$ will also have related symmetries, and thus can be reduced to the equations of motion of the Toda theory associated to $\hat{g}_2$ by identifying certain fields. Also, solutions to the $\hat{g}_1$ theory may be reduced (“folded”) to solutions to the $\hat{g}_2$ theory [6, 16]. Furthermore, every solution to the $\hat{g}_2$ theory can be unfolded to a solution to the $\hat{g}_1$ theory, so we only need to deal with simply-laced loop algebras.

Finally, we want to point out how plane-wave solutions come out of the formalism. If we replace the single-soliton ansatz (2.13) by the plane-wave ansatz

$$\tau_j^{(n)} = \delta_j^{(n)} e^{in\Phi}, \quad (2.28)$$

where $\Phi = (kx - \omega t) + \xi$, and $\omega^2 - k^2 = m^2\lambda$, most of the discussion on the solitons is still valid for the plane waves. Even though we will only write down explicitly the soliton—soliton

scattering solutions, soliton—plane-wave and wave-wave scattering solutions can be easily derived by changing some parameters from real to purely imaginary.

3 $\hat{A}_r$ theory

The Dynkin diagram for $A_r^{(1)}$ is shown in Figure 3.1.




Figure 3.1: Affine Dynkin diagram for $\hat{A}_r$.

Since $l_j = 1$, for all j 's, so $L_{jk} = K_{jk}$, and Hirota's equation (2.5) reads

$$\frac{1}{2m^2}(D_t^2 - D_x^2) \tau_j \cdot \tau_j = (\tau_{j+1}\tau_{j-1} - \tau_j^2), \quad \text{for } j = 0, 1, 2, \dots, r. \quad (3.1)$$

Due to the periodicity of the extended Dynkin diagram, it is implied that $\tau_{j+r+1} = \tau_j$. The explicit expression for $B_j(k_1, \dots, k_N)$ is given below:

$$B_j(k_1, \dots, k_N) = \sum_{0 \prec (m_1, \dots, m_N) \prec (k_1, \dots, k_N)} \delta_{j+1}(m_1, \dots, m_N) \delta_{j-1}(k_1 - m_1, \dots, k_N - m_N). \quad (3.2)$$

The matrix L has eigenvalues and eigenvectors

$$\lambda_s = 4 \sin^2 \vartheta_s, \quad \Lambda_s = (1, e^{2i\vartheta_s}, e^{4i\vartheta_s}, \dots, e^{2ir\vartheta_s}), \quad \text{for } s=1, 2, \dots, r, \quad (3.3)$$

where $\vartheta_s = s\pi/h$, and $h = r + 1$ is the Coxeter number.

For any given integer $1 \leq s \leq r$, there is a class of single-soliton solutions ($\kappa = 1$) given in [1], with the only non-vanishing coefficients $\delta^{(1)} = \Lambda_s$. The topological charge of the soliton depends on the value of the parameter ξ . Because of degeneracy in eigenvalues, $\lambda_s = \lambda_{h-s}$, there also exist $\kappa = 2$ solitons [7]:

$$\delta_j^{(1)} = y^+ \exp(2\pi i s j / h) + y^- \exp(-2\pi i s j / h), \quad \delta_{s,j}^{(2)} = y^+ y^- \cos^2 \vartheta_s. \quad (3.4)$$

Here y^+ and y^- are independent variables, if we take either of y^+ and y^- be zero in the above equations, we will get back to $\kappa = 1$ solutions. In the following we will construct multi-soliton solutions consisting of solitons with $\kappa = 2$, since if there are some $\kappa = 1$ solitons involved, we only need to take some of the y 's to be zero.

First we solve for a two-soliton solution. Let $\lambda^{(1)} = \lambda_p$, $\lambda^{(2)} = \lambda_q$, p, q are integers with $1 \leq p, q \leq r$, and $\kappa_1 = \kappa_2 = 2$, from (2.15), (2.25) and (3.4),

$$\begin{aligned}\delta_j(1, 0) &= y_1^+ \exp(2\pi i p j / h) + y_1^- \exp(-2\pi i p j / h), \\ \delta_j(0, 1) &= y_2^+ \exp(2\pi i q j / h) + y_2^- \exp(-2\pi i q j / h), \\ \delta_j(2, 0) &= y_1^+ y_1^- \cos^2 \vartheta_p, \quad \delta_j(0, 2) = y_2^+ y_2^- \cos^2 \vartheta_q.\end{aligned}\tag{3.5}$$

By straightforward calculations, we get:

$$\begin{aligned}\delta_j(1, 1) &= y_1^+ y_2^+ e^{\gamma_{pq}^{++}} \exp(2\pi i (p + q) j / h) + y_1^- y_2^- e^{\gamma_{pq}^{--}} \exp(-2\pi i (p + q) j / h) \\ &\quad + y_1^+ y_2^- e^{\gamma_{pq}^{+-}} \exp(2\pi i (p - q) j / h) + y_1^- y_2^+ e^{\gamma_{pq}^{-+}} \exp(-2\pi i (p - q) j / h), \\ \delta_j(2, 1) &= \delta_j(2, 0) \delta_j(0, 1) \exp(\gamma_{pq}^{++} + \gamma_{pq}^{+-}), \\ \delta_j(1, 2) &= \delta_j(0, 2) \delta_j(1, 0) \exp(\gamma_{pq}^{++} + \gamma_{pq}^{+-}), \\ \delta_j(2, 2) &= \delta_j(2, 0) \delta_j(0, 2) \exp(2\gamma_{pq}^{++} + 2\gamma_{pq}^{+-}),\end{aligned}\tag{3.6}$$

and all higher order terms vanish. Here

$$\exp(\gamma_{s_a, s_b}^{++}) = \exp(\gamma_{s_a, s_b}^{--}) = \exp(\gamma_{s_a, s_b}^+) = \frac{\cosh \theta_{ab} - \cos \vartheta_{s_a - s_b}}{\cosh \theta_{ab} - \cos \vartheta_{s_a + s_b}},\tag{3.7}$$

$$\exp(\gamma_{s_a, s_b}^{+-}) = \exp(\gamma_{s_a, s_b}^{-+}) = \exp(\gamma_{s_a, s_b}^-) = \frac{\cosh \theta_{ab} + \cos \vartheta_{s_a + s_b}}{\cosh \theta_{ab} + \cos \vartheta_{s_a - s_b}}.\tag{3.8}$$

Constructing an N -soliton solution means finding all nonvanishing $\delta_j(m_1, \dots, m_N)$. From (2.25), if any of the m_a vanishes, then the corresponding δ can be taken from some $N - 1$ soliton solutions. Since we have assumed $\kappa = 2$, from (2.26), if any of the m_a is greater than or equal to 3, then the corresponding δ shall vanish. So we only need to consider

$1 \leq m_a \leq 2$. Furthermore from the above two-soliton solutions we see that $\delta_j(2, 1)$ and $\delta_j(1, 2)$ are related by renaming the solitons, or shuffling the indices $p \leftrightarrow q$. This is also true for N-soliton solutions, so we only have to find $\delta_j(\overbrace{2, \dots, 2}^i, \overbrace{1, \dots, 1}^{N-i})$, for $i = 0, 1, \dots, N$.

Let $\lambda^{(a)} = \lambda_{s_a}$, with $1 \leq s_a \leq r$, We list the nontrivial coefficients for N-soliton solutions below:

$$\delta_j(\overbrace{1, \dots, 1}^N) = \sum_{\{\sigma\}} y_1^{\sigma_1} \cdots y_N^{\sigma_N} \exp(i2\pi(\sigma_1 s_1 + \cdots + \sigma_N s_N)j/h) \exp\left(\sum_{1 \leq b < c \leq N} \gamma_{s_b, s_c}^{\sigma_b \sigma_c}\right), \quad (3.9)$$

$$\delta_j(\overbrace{2, \dots, 2}^i, \overbrace{1, \dots, 1}^{N-i}) = \delta_j(\overbrace{2, \dots, 2}^i, \overbrace{0, \dots, 0}^{N-i}) \delta_j(\overbrace{0, \dots, 0}^i, \overbrace{1, \dots, 1}^{N-i}) \prod_{\substack{a \leq i \\ b > i}} \exp(\gamma_{s_a, s_b}^+ + \gamma_{s_a, s_b}^-), \quad (3.10)$$

$$\delta_j(\overbrace{2, \dots, 2}^N) = \delta_{1,j}^{(2)} \cdots \delta_{N,j}^{(2)} \prod_{a < b} \exp(2\gamma_{s_a, s_b}^+ + 2\gamma_{s_a, s_b}^-). \quad (3.11)$$

We will outline our construction and notation in three steps, assuming all the $N - 1$ soliton solutions are known.

Step 1: To get (3.9).

One can easily see that $\delta_j(\overbrace{1, \dots, 1}^N)$ should be N-linear in the y 's. Note that $y_1^+ y_1^-$ is considered to be quadratic while $y_1^+ y_2^-$ is bilinear. So

$$\delta_j(\overbrace{1, \dots, 1}^N) = \sum_{\{\sigma\}} C(\sigma_1, \dots, \sigma_N) y_1^{\sigma_1} \cdots y_N^{\sigma_N},$$

where $\sigma_a = \pm 1$ and $C(\sigma_1, \dots, \sigma_N)$ are independent of x, t and all the y 's. y^σ should be understood as y^+ or y^- , Summation is over all possible combinations of σ 's

If we take $y_1^- = \cdots = y_N^- = 0$, then only one term survives, and all the κ 's become 1 instead of 2, so we should get those of Hollowood,¹ i.e.,

$$C(1, \dots, 1) = \exp\left(\sum_{1 \leq b < c \leq s_1} \gamma_{s_b, s_c}^+\right) \exp(i2\pi(s_1 + \cdots + s_N)j/h). \quad (3.12)$$

¹In [1] the N-soliton solutions were written down without explicit proof.

Proving the above equation is equivalent to proving the following identity:

$$\sum_{\{\sigma\}} \left(f(\sigma_1, \sigma_2, \dots, \sigma_N) - 4 \sin^2(\sigma_1 \vartheta_1 + \dots + \sigma_N \vartheta_N) \right) \prod_{a < b} \left(\cosh \theta_{ab} - \cos(\sigma_a \vartheta_a - \sigma_b \vartheta_b) \right) = 0, \quad (3.13)$$

where each σ takes value 1 or -1 , and summation is over all possible combinations of σ .

Since directly proving the above identity is hard, we will construct (3.12) without direct proof of (3.13), by using (2.27) instead. Since we have taken $y_1^- = \dots = y_N^- = 0$, all the single solitons have $\kappa = 1$. By (2.26), the only nonvanishing terms in $V_j(1, \overbrace{2, \dots, 2}^{N-1})$ are

$$\begin{aligned} & \delta_{j-1}(1, 1, \dots, 1) \delta_{j+1}(0, 1, \dots, 1) + \delta_{j+1}(1, 1, \dots, 1) \delta_{j-1}(0, 1, \dots, 1) \\ & - 2 \cos(2\pi s_1/h) \delta_j(1, 1, \dots, 1) \delta_j(0, 1, \dots, 1). \end{aligned}$$

Let $\delta_j(1, 1, \dots, 1) = e^{i2\pi j(s_2 + \dots + s_N)/h} T_j$, from the vanishing condition of $V_j(1, \overbrace{2, \dots, 2}^{N-1})$ we get:

$$T_{j-1} + T_{j+1} - 2 \cos(2\pi s_1/h) T_j = 0. \quad (3.14)$$

The above homogeneous second order difference equation has two linearly independent solutions:

$$T_j = T e^{\pm i2\pi j s_1/h}.$$

From a symmetry argument or the vanishing conditions $V_j(2, \dots, 2, 1) = 0$, we can see that we should chose the “+” sign in the above solution. The constant factor T can be determined from the vanishing condition of $V_j(1, 1, \overbrace{2, \dots, 2}^{N-2})$, which gives (3.12). We could find all the $C(\sigma_1, \dots, \sigma_N)$ ’s the same way, thus proving (3.9).

Step 2: To get (3.10).

First let $y_{i+1}^- = \dots = y_N^- = 0$, so $\kappa_1 = \dots = \kappa_i = 2$, $\kappa_{i+1} = \dots = \kappa_N = 1$. Adding all the nonvanishing terms in $V_j(\overbrace{4, \dots, 4}^i, \overbrace{2, \dots, 2}^{N-i-1}, 1)$, from Eq. (2.27), the sum should vanish. So we obtain a homogeneous second order difference equation of $\delta_j(\overbrace{2, \dots, 2}^i, 1, \dots, 1)$. We can write down the vanishing condition of $V_j(\overbrace{0, \dots, 0}^i, \overbrace{2, \dots, 2}^{N-i-1}, 1)$ also, which gives a

homogeneous second order difference equation of $\delta_j(\overbrace{0, \dots, 0}^i, 1, \dots, 1)$. Comparing the two equations, and from the fact that they obey the same boundary conditions and have similar symmetries, we see that $\delta_j(\overbrace{2, \dots, 2}^i, 1, \dots, 1)$ should be proportional to $\delta_j(\overbrace{0, \dots, 0}^i, 1, \dots, 1)$.

The proportionality constant can be determined from the vanishing condition:

$$V_j(\overbrace{4, \dots, 4}^i, \overbrace{2, \dots, 2}^{N-i-2}, 1, 1) = 0.$$

Second, there should be 2^{N-i} terms, and each can be derived in a similar way. Putting them all together, we will get (3.10).

Step 3: To get (3.11).

From the vanishing condition of $V(4, \dots, 4, 3)$ and using factorization of (3.10), we obtain:

$$(\lambda_N - 2)\delta_j(2, \dots, 2) + e^{\pm i 2\pi s_N/h} \delta_{j-1}(2, \dots, 2) + e^{\mp i 2\pi s_N/h} \delta_{j+1}(2, \dots, 2) = 0. \quad (3.15)$$

Adding and subtracting the two equations, we see that $\delta_j(2, \dots, 2)$ is constant, which constant can be calculated from the vanishing conditions of $V(4, \dots, 4, 3, 3)$.

This completes the construction. These $\kappa = 2$ solitons are not really “single solitons” in the strict sense, since they change shape after scattering with other ($\kappa = 1$) solitons. In general a $\kappa = 2$ soliton has two peaks, and each peak behaves as if it is a free soliton when scattering off other solitons. These $\kappa = 2$ solitons should be interpreted as compounds of two $\kappa = 1$ solitons interacting in a peculiar way so that there seems to be no interaction among the peaks. Indeed, one can obtain such $\kappa = 2$ soliton solutions by taking the static limit of some two-soliton scattering solutions of $\kappa = 1$ solitons [18].

Let us look into the scattering of two $\kappa = 1$ solitons in a little more detail. Since the only effect after scattering is a shift in the center of mass for each soliton, and the shift does not depend directly on the topological charge of each soliton but rather on the topological classes they belong to, one may be tempted to believe that the interactions among the solitons

depend only on the classes. Furthermore, since γ_{ab}^+ in (3.7) is always negative, it appears as if the interactions are always attractive. However, on closer inspection, one notices that the two solitons could either speed up or slow down as they approach each other. If they speed up, eventually the two peaks will merge together and then separate again. If, on the other hand, they slow down, as they approach to within a certain distance, they begin to exchange shape, topological charge and other attributes, and then move apart again as if the two solitons have switched position. Because the two solitons performed this exchange over some finite distance, even though the interaction is repulsive, the total shift looks like an attractive case. In both cases, there is no reflection even if the masses are equal. Details of which pair of solitons attract or repel each other still need to be worked out. Since we can also have two solitons stand still as if there is no interaction at all, the interaction cannot be described by a simple equivalent potential. This may be because the solitons are extended objects.

If $r = 2N - 1$, for some integer $N = 2, 3, \dots$, and

$$\tau_j = \tau_{r-j}, \quad \text{for } j = 1, 2, \dots, N, \quad (3.16)$$

then the set $\tau_0, \tau_1, \dots, \tau_N$ will form a tau-function solution set for $\text{sp}(2N)$ affine Toda models. It is easy to check that if we set $y^+ = y^-$, then the $\kappa = 2$ soliton given in (3.4) will satisfy the condition (3.16). These $\kappa = 2$ solitons of $\text{sp}(2N)$ do look and behave like single solitons: the two overlapping peaks do not drift apart since they are protected by the symmetry (3.16). From the solutions (3.9) to (3.11) one can also easily obtain multi-soliton solutions for $\text{sp}(2N)$ affine TFTs.

4 $\hat{D}_4$

Dynkin diagram for $\hat{D}_4$ is shown below:




Figure 5.1: Affine Dynkin diagram for $\hat{D}_4$.

$$l_0 = l_1 = l_3 = l_4 = 1, \quad l_2 = 2.$$

$$B_0(k_1, \dots, k_N) = B_1(k_1, \dots, k_N) = B_3(k_1, \dots, k_N) = B_4(k_1, \dots, k_N) = 0,$$

$$B_2(k_1, \dots, k_N) = \sum \delta_0(m_1, \dots, m_N) \delta_1(n_1, \dots, n_N) \delta_3(p_1, \dots, p_N) \delta_4(q_1, \dots, q_N),$$

where the summation is over $p_a + q_a + m_a + n_a = k_a$. The eigenvalues and eigenvectors of the matrix L_{jk} are:

$$\lambda_1 = 2; \quad \Lambda_1 = (1, -1, 0, 1, -1),$$

$$\lambda_2 = 2; \quad \Lambda_2 = (1, 1, 0, -1, -1),$$

$$\lambda_3 = 2; \quad \Lambda_3 = (1, -1, 0, -1, 1),$$

$$\lambda_4 = 6; \quad \Lambda_4 = (1, 1, -4, 1, 1),$$

We also define $\Lambda_6 \equiv (0, 0, 1, 0, 0)$ for later convenience.

For $\lambda = 6$, the eigenvalue is not degenerate, there is only one corresponding single soliton solution with $\kappa = 1$:

$$\delta^{(1)} = \Lambda_4; \quad \delta^{(2)} = \Lambda_6.$$

However, the eigenvalue $\lambda = 2$ has degeneracy 3, so one can construct a $\kappa = 3$ soliton:

$$\begin{aligned} \delta^{(1)} &= y_1 \Lambda_1 + y_2 \Lambda_2 + y_3 \Lambda_3, \\ \delta^{(2)} &= \frac{1}{3} (y_1 y_2 \Lambda_3 + y_2 y_3 \Lambda_1 + y_1 y_3 \Lambda_2) + (y_1^2 + y_2^2 + y_3^2) \Lambda_6, \\ \delta^{(3)} &= P_3 (\Lambda_4 - 12 \Lambda_6), \quad \delta^{(4)} = P_4 \Lambda_6, \\ \delta^{(5)} &= 0, \quad \delta^{(6)} = P_3^2 \Lambda_6, \end{aligned} \tag{4.17}$$

where $P_3 = y_1 y_2 y_3 / 27$, $P_4 = y_1^2 y_2^2 + y_1^2 y_3^2 + y_2^2 y_3^2 / 9$. One can obtain three $\kappa = 2$ or $\kappa = 1$ solitons if one or two of the y 's is taken to be zero.

We will first study the two-soliton scattering solutions. Denote:

$$C_1 = \frac{\cosh \theta - 1}{\cosh \theta + 1}, \quad C_2 = \frac{2 \cosh \theta - 1}{2 \cosh \theta + 1}, \quad C_3 = \frac{2 \cosh \theta - \sqrt{3}}{2 \cosh \theta + \sqrt{3}}, \quad C_4 = \frac{2 \cosh \theta + 1}{2 \cosh \theta - 1}. \quad (4.18)$$

Since it is straight forward to calculate the double-soliton solutions, we will write down the nontrivial coefficients without derivation.

For $\lambda^{(1)} = \lambda^{(2)} = 6$:

$$\begin{aligned} \delta(1, 0) &= \delta(0, 1) = \Lambda_4, & \delta(2, 0) &= \delta(0, 2) = \Lambda_6, \\ \delta(1, 1) &= \frac{2(5 \cosh \theta - 4)}{3(\cosh \theta + 1)} \Lambda_0 + \frac{(11 - 14 \cosh \theta)}{3(2 \cosh \theta + 1)} \Lambda_4, \\ \delta(1, 2) &= \delta(2, 1) = -4C_1 C_2 \Lambda_6, & \delta(2, 2) &= C_1^2 C_2^2 \Lambda_6. \end{aligned} \quad (4.19)$$

For $\lambda^{(1)} = 2$ with $\kappa = 3$, and $\lambda^{(2)} = 6$ with $\kappa = 1$: $\delta(m, 0)$ will be taken as $\delta^{(m)}$ in (4.17), $\delta(0, 1) = \Lambda_4$, $\delta(0, 2) = \Lambda_6$, and

$$\begin{aligned} \delta(1, 1) &= C_3(y_1 \Lambda_1 + y_2 \Lambda_3 + y_3 \Lambda_3), \\ \delta(2, 1) &= \frac{1}{3} C_3^2 (y_1 y_2 \Lambda_3 + y_2 y_3 \Lambda_1 + y_1 y_3 \Lambda_2) - 4C_3(y_1^2 + y_2^2 + y_3^2) \Lambda_6, \\ \delta(2, 2) &= C_3^2 (y_1^2 + y_2^2 + y_3^2) \Lambda_6, \\ \delta(3, 1) &= P_3 C_3^3 \Lambda_4 + 4P_3(9C_3^2 + 9C_3 - 1) \Lambda_6, \\ \delta(3, 2) &= 16P_3 C_3^3 \Lambda_6, \\ \delta(4, 1) &= -4P_4 C_3^2 \Lambda_6, & \delta(4, 2) &= P_4 C_3^4 \Lambda_6, \\ \delta(6, 1) &= -4P_3^2 C_3^3 \Lambda_6, & \delta(6, 2) &= P_3^2 C_3^6 \Lambda_6. \end{aligned} \quad (4.20)$$

For $\lambda^{(1)} = \lambda^{(2)} = 2$ with $\kappa_1 = 3$, $\kappa_2 = 1$: Let $\delta(m, 0)$ will be taken as $\delta^{(m)}$ in (4.17), $\delta(0, 1) = \Lambda_1$, $\delta(0, 2) = \Lambda_6$, and

$$\delta(1, 1) = y_1 \left(C_1 C_4 \Lambda_0 - \frac{2(2 \cosh \theta - 3)}{2 \cosh \theta - 1} \Lambda_6 \right) + C_2 (y_2 \Lambda_3 + y_3 \Lambda_2),$$

$$\begin{aligned}
\delta(2, 1) &= y_2 y_3 \left(\frac{1}{3} C_2^2 \Lambda_4 + \frac{4(2 \cosh \theta - 3)}{3(2 \cosh \theta + 1)} \Lambda_6 \right) + \frac{1}{3} C_1 y_1 (y_2 \Lambda_2 + y_3 \Lambda_3), \\
\delta(2, 2) &= (y_1^2 C_1^2 C_4^2 + y_2^2 C_2^2 + y_3^2 C_2^2) \Lambda_6, \\
\delta(3, 1) &= \frac{1}{27} y_1 y_2 y_3 C_1 C_2 \Lambda_1 + \frac{4y_1(y_2^2 + y_3^2)}{3(1 + \cosh \theta)(1 + 2 \cosh \theta)} \Lambda_6, \\
\delta(3, 2) &= -\frac{16}{27} C_1 C_2 y_1 y_2 y_3 \Lambda_6, \\
\delta(4, 1) &= \frac{16}{27} \frac{C_1}{(1 - 2 \cosh \theta)(1 + 2 \cosh \theta)} y_1 y_2 y_3 \Lambda_6, \\
\delta(4, 2) &= \frac{1}{9} (C_1^2 y_1^2 (y_2^2 + y_3^2) + C_2^4 y_2^2 y_3^2) \Lambda_6, \\
\delta(5, 1) &= \frac{4}{81} \frac{C_2 y_1 y_2^2 y_3^2}{(1 + \cosh \theta)(1 + 2 \cosh \theta)} \Lambda_6, \\
\delta(6, 2) &= \frac{1}{729} C_1^2 C_2^2 y_1^2 y_2^2 y_3^2 \Lambda_6.
\end{aligned} \tag{4.21}$$

From the solutions (4.19) and (4.20), with a little analysis, we can see that the solitons retain their shape after scattering. However in the solution (4.21), the $\kappa = 3$ soliton changes shape after scattering, as signified by the fact that the constant y_1 is treated differently from y_2 and y_3 . The $\kappa = 3$ soliton can be thought of as a composite of three $\kappa = 1$ solitons. We will substantiate this claim by writing down the three-soliton scattering solution. Let $\delta(1, 0, 0) = y_1 \Lambda_1$, $\delta(0, 1, 0) = y_2 \Lambda_2$, $\delta(0, 0, 1) = y_3 \Lambda_3$. Then $\delta(2, 0, 0) = y_1^2 \Lambda_6$, $\delta(0, 2, 0) = y_2^2 \Lambda_6$, $\delta(0, 0, 2) = y_3^2 \Lambda_6$, and

$$\begin{aligned}
\delta(1, 1, 0) &= y_1 y_2 D_{12} \Lambda_3, \quad \delta(1, 0, 1) = y_1 y_3 D_{13} \Lambda_2, \quad \delta(0, 1, 1) = y_2 y_3 D_{23} \Lambda_1, \\
\delta(1, 1, 1) &= y_1 y_2 y_3 D_{12} D_{23} D_{13} \Lambda_0 - y_1 y_2 y_3 (2D_{12} D_{23} D_{13} + 16D_{123}) \Lambda_6, \\
\delta(2, 2, 0) &= y_1^2 y_2^2 D_{12}^2 \Lambda_6, \quad \delta(2, 0, 2) = y_1^2 y_3^2 D_{13}^2 \Lambda_6, \quad \delta(0, 2, 2) = y_2^2 y_3^2 D_{23}^2 \Lambda_6, \\
\delta(2, 2, 2) &= y_1^2 y_2^2 y_3^2 D_{12}^2 D_{23}^2 D_{13}^2 \Lambda_6,
\end{aligned} \tag{4.22}$$

where $D_{ab} = \frac{2 \cosh \theta_{ab} - 1}{2 \cosh \theta_{ab} + 1}$ and $D_{123} = \frac{1}{(2 \cosh \theta_{12} + 1)(2 \cosh \theta_{23} + 1)(2 \cosh \theta_{13} + 1)}$. Now if all the three-solitons have the same velocity, then $\theta_{12} = \theta_{23} = \theta_{13} = 0$, and we can have $\Gamma_1 = \Gamma_2 = \Gamma_3 = \Gamma$. Then the above three-soliton scattering solution will have the single soliton form with $\delta^{(m)} = \sum_{m_1+m_2+m_3=m} \delta(m_1, m_2, m_3)$, which gives the solution set (4.17).

Because of the non-trivial symmetries of the Dynkin diagram, the solutions can be folded to yield solutions to G_2 affine TFT by demanding $\tau_1 = \tau_3 = \tau_4$. The $\kappa = 3$ soliton with $y_1 = y_2 = y_3$ and the $\lambda = 6$ soliton survive the folding procedure. Protected by the discrete symmetry, these will be true single solitons of G_2 affine TFT.

5 $D_{2r}^{(1)}$ Theory for $r \geq 5$

The Dynkin diagram for $D_r^{(1)}$ ($r \geq 5$) is shown in Figure 4.1.




Figure 4.1: Affine Dynkin diagram for $D_r^{(1)}$.

$l_0 = l_1 = l_{r-1} = l_r = 1$, $l_j = 2$ for $2 \leq j \leq r-2$. The explicit form of $B_j(k_1, \dots, k_N)$ is given below:

$$\begin{aligned}
B_0(k_1, \dots, k_N) &= B_1(k_1, \dots, k_N) = B_{r-1}(k_1, \dots, k_N) = B_r(k_1, \dots, k_N) = 0, \\
B_2(k_1, \dots, k_N) &= \sum_{p_a + q_a + m_a = k_a} \delta_0(p_1, \dots, p_N) \delta_1(q_1, \dots, q_N) \delta_3(m_1, \dots, m_N), \\
B_{r-2}(k_1, \dots, k_N) &= \sum_{p_a + q_a + m_a = k_a} \delta_r(p_1, \dots, p_N) \delta_{r-1}(q_1, \dots, q_N) \delta_{r-3}(m_1, \dots, m_N),
\end{aligned} \tag{5.1}$$

and the rest of the $B_j(k_1, \dots, k_N)$'s are the same as in (3.2).

The matrix L has eigenvalues and eigenvectors

$$\begin{aligned}
\lambda_s &= 8 \sin^2 \vartheta_s, \quad \text{for } 1 \leq s \leq r-2; \\
\Lambda_s &= \left(1, 1, 2 \frac{\cos 3\vartheta_s}{\cos \vartheta_s}, 2 \frac{\cos 5\vartheta_s}{\cos \vartheta_s}, \dots, 2 \frac{\cos (2r-5)\vartheta_s}{\cos \vartheta_s}, (-1)^s, (-1)^s \right), \\
\lambda_{r-1} &= \lambda_r = 2; \quad \Lambda_{r-1} = (1, -1, 0, 0, \dots, 0), \quad \Lambda_r = (0, 0, \dots, 0, -1, 1),
\end{aligned} \tag{5.2}$$

where $\vartheta_s = s\pi/2(r-1) = s\pi/h$.

First we establish some useful identities for later calculations of double-soliton solutions:

$$\sum_{j=2}^{r-2} \cos(2j-1)\vartheta_p = \begin{cases} r-3, & \text{if } p=0 \bmod h; \\ -(1+(-1)^p) \cos \vartheta_p, & \text{otherwise} \end{cases} \quad (5.3)$$

$$\sum_{j=2}^{r-2} \frac{\cos(2j-1)\vartheta_p \cos(2j-1)\vartheta_q}{\cos \vartheta_p \cos \vartheta_q} = \begin{cases} -1 + \frac{r-1}{2 \cos^2 \vartheta_p}, & \text{if } p \pm q = 0 \bmod h; \\ -(1+(-1)^{p+q}), & \text{otherwise} \end{cases} \quad (5.4)$$

$$\begin{aligned} & \sum_{j=2}^{r-2} \frac{\cos(2j-1)\vartheta_p \cos(2j-1)\vartheta_q \cos(2j-1)\vartheta_m}{\cos \vartheta_p \cos \vartheta_q \cos \vartheta_m} \\ &= \begin{cases} -2 + \frac{r-1}{4 \cos \vartheta_p \cos \vartheta_q \cos \vartheta_m}, & \text{if } p \pm q \pm m = 0 \bmod h; \\ -(1+(-1)^{p+q+m}), & \text{otherwise} \end{cases} \end{aligned}$$

$$\begin{aligned} & \sum_{j=2}^{r-2} \frac{(\cos(2j-3)\vartheta_p \cos(2j+1)\vartheta_q + \cos(2j+1)\vartheta_p \cos(2j-3)\vartheta_q) \cos(2j-1)\vartheta_m}{\cos \vartheta_p \cos \vartheta_q \cos \vartheta_m} \\ &= \begin{cases} -2 \left(\frac{\cos 3\vartheta_p}{\cos \vartheta_p} + \frac{\cos 3\vartheta_p}{\cos \vartheta_p} \right) + \frac{(r-1) \cos 2\vartheta_{p+q}}{\cos \vartheta_p \cos \vartheta_q \cos \vartheta_m}, & \text{if } m \pm (p-q) = 0 \bmod h; \\ -2 \left(\frac{\cos 3\vartheta_p}{\cos \vartheta_p} + \frac{\cos 3\vartheta_p}{\cos \vartheta_p} \right) + \frac{(r-1) \cos 2\vartheta_{p-q}}{\cos \vartheta_p \cos \vartheta_q \cos \vartheta_m}, & \text{if } m \pm (p+q) = 0 \bmod h; \\ -(1+(-1)^{p+q+m}) \left(\frac{\cos 3\vartheta_p}{\cos \vartheta_p} + \frac{\cos 3\vartheta_p}{\cos \vartheta_p} \right), & \text{otherwise.} \end{cases} \end{aligned}$$

Single-soliton solutions have been given in [7]. For $1 \leq s \leq r-2$, the eigenvalues are non-degenerate, $\kappa = 1$,

$$\delta^{(1)} = \Lambda_s, \quad \delta^{(2)} = u \equiv (0, 0, 1, \dots, 1, 0, 0). \quad (5.5)$$

For the eigenvalue $\lambda = 2$, which is doubly degenerate, we can have a $\kappa = 2$ soliton:

$$\begin{aligned} \delta^{(1)} &= y_1 \Lambda_{r-1} + y_2 \Lambda_r, \\ \delta^{(2)} &= \left(c, c, d_2, d_3, \dots, d_{r-2}, (-1)^{r-1}c, (-1)^{r-1}c \right), \\ \delta^{(4)} &= c^2 u = c^2 (0, 0, 1, 1, \dots, 1, 0, 0), \end{aligned} \quad (5.6)$$

where $c = (y_1^2 + (-1)^{r-1}y_2^2)/2h$, and $d_j = (-1)^j((h-2j+1)y_1^2 + (-1)^r(2j-1)y_2^2)/h$. If we let $y_2 = \pm(-1)^{r/2}y_1$, then the above $\kappa = 2$ soliton will be reduced to a $\kappa = 1$ soliton.

We will next write down some double-soliton solutions without derivation. Let $\lambda^{(1)} = \lambda_p$, and $\lambda^{(2)} = \lambda_q$, for some integers $1 \leq p, q \leq r-2$, then $\delta_j(1, 0) = \Lambda_{p,j}$, $\delta_j(0, 1) = \Lambda_{q,j}$, and

$$\begin{aligned}
\delta_j(0, 2) &= \delta_j(2, 0) = u_j, \\
\delta_0(1, 1) &= \delta_1(1, 1) = -1 + \frac{\cos(\vartheta_p + \vartheta_q)}{\cos \vartheta_p \cos \vartheta_q} e^{\gamma_{pq}^+} + \frac{\cos(\vartheta_p - \vartheta_q)}{\cos \vartheta_p \cos \vartheta_q} e^{\gamma_{pq}^-}, \\
\delta_r(1, 1) &= \delta_{r-1}(1, 1) = (-1)^{p+q} \delta_0(1, 1), \\
\delta_j(1, 1) &= \frac{2 \cos(2j-1) \vartheta_{p+q}}{\cos \vartheta_p \cos \vartheta_q} e^{\gamma_{pq}^+} + \frac{2 \cos(2j-1) \vartheta_{p-q}}{\cos \vartheta_p \cos \vartheta_q} e^{\gamma_{pq}^-}, \quad j=2, \dots, r-2, \quad (5.7) \\
\delta_j(2, 1) &= u_j \delta_j(1, 0) \delta_0(1, 1), \\
\delta_j(1, 2) &= u_j \delta_j(0, 1) \delta_0(1, 1), \\
\delta_j(2, 2) &= u_j \left(\delta_0(1, 1) \right)^2.
\end{aligned}$$

Let $\lambda^{(2)} = \lambda_p$ for some $0 < p \leq r-2$, $\lambda^{(2)} = 2$, and $\delta_j(1, 0) = \Lambda_{p,j}$, $\delta_j(0, 1) = \Lambda_{r-1} \pm (-1)^{r/2} \Lambda_r$, then:

$$\begin{aligned}
\delta(2, 0) &= u, \quad \delta(0, 2) = \left(0, 0, 1, -1, 1, -1, \dots, (-1)^{r-2}, 0, 0 \right), \\
\delta(1, 1) &= (\Lambda_{r-1} \pm (-1)^{\frac{r}{2}+p} \Lambda_r) e^{\tilde{\gamma}}, \\
\delta_j(1, 2) &= (-1)^j \Lambda_{p,j} e^{\tilde{\gamma}}, \quad \text{for } 2 \leq j \leq r-2, \quad (5.8) \\
\delta_j(2, 2) &= \delta_j(0, 2) e^{2\tilde{\gamma}},
\end{aligned}$$

where $e^{\tilde{\gamma}} = (\cosh \theta - \sin \vartheta_p) / (\cosh \theta + \sin \vartheta_p)$.

Let $\lambda^{(1)} = \lambda^{(2)} = 2$, $\delta(1, 0) = z_1(\Lambda_{r-1} + (-1)^{r/2} \Lambda_r)$ and $\delta(0, 1) = z_1(\Lambda_{r-1} - (-1)^{r/2} \Lambda_r)$, then:

$$\begin{aligned}
\delta(2, 0)/z_1^2 &= \delta(0, 2)/z_2^2 = (0, 0, 1, -1, 1, -1, \dots, (-1)^{r-1}, 0, 0) \\
\delta(1, 1) &= z_1 z_2 \frac{1 + (-1)^r}{r-1} \frac{\cosh \theta}{\cosh \theta + 1} \Lambda_0 \\
&\quad + z_1 z_2 \sum_{s=1}^{r-2} \frac{2 \cos^2 \vartheta_s \left(1 + (-1)^{(r+s+1)} \right)}{r-1} \frac{\cosh \theta - 2 \sin^2 \theta_s}{\cosh \theta - 2 \sin^2 \vartheta + 1} \Lambda_s \quad (5.9) \\
\delta(2, 2) &= \delta_0^2(1, 1) u.
\end{aligned}$$

Taking the static limit of the above double-soliton solution, and letting $y_1 = z_1 + z_2$ and $y_2 = (-1)^{r/2}(z_1 - z_2)$, it will then have single-soliton form with $\delta^{(m)} = \sum_{m_1+m_2=m} \delta(m_1, m_2)$, which gives the $\kappa = 2$ solution of (5.6).

We are interested in presenting an N -soliton solution. As in section 3, it will be built on some $(N - 1)$ -soliton solutions, and for simplicity, we assume that all solitons involved are of the form (5.5). Here we list all the non-trivial coefficients:

$$\delta_0(1, 1, \dots, 1) = \delta_1(1, 1, \dots, 1) = \prod_{a < b} \delta_{ab,0}(1, 1), \quad (5.10)$$

$$\delta_r(1, 1, \dots, 1) = \delta_{r-1}(1, 1, \dots, 1) = (-1)^{s_1+s_2+\dots+s_N} \delta_0(1, 1, \dots, 1), \quad (5.11)$$

$$\delta_j(1, 1, \dots, 1) = \sum_{\{\sigma\}} \frac{\cos((2j-1)(\sigma_1\vartheta_{s_1} + \sigma_2\vartheta_{s_2} + \dots + \sigma_N\vartheta_{s_N}))}{\cos\vartheta_{s_1} \cos\vartheta_{s_2} \dots \cos\vartheta_{s_N}} \prod_{a < b} e^{\gamma_{ab}^{\sigma_a \sigma_b}}, \quad (5.12)$$

$$\delta_j(\overbrace{2, \dots, 2}^i, 1, \dots, 1) = \delta_j(2, \dots, 2, 0, \dots, 0) \delta_j(0, \dots, 0, 1, \dots, 1) \prod_{\substack{a \leq i \\ b > i}} \delta_{ab,0}(1, 1), \quad (5.13)$$

$$\delta_j(2, \dots, 2) = \delta_j^{(2)} \delta_0^2(1, \dots, 1). \quad (5.14)$$

In Eqs. (5.12), (5.13) and (5.14), j runs from 2 to $r-2$. We sketch our construction below.

One can easily write down the second-order difference equation for $\delta_j(2, \dots, 2)$ from the vanishing conditions of $V_j(4, \dots, 4)$:

$$\delta_j^2(2, \dots, 2) - \delta_{j-1}(2, \dots, 2) \delta_{j+1}(2, \dots, 2) = 0, \quad \text{for } j = 3, 4, \dots, r-3. \quad (5.15)$$

We need two initial conditions to solve (5.15):

$$\delta_0(2, \dots, 2) = 0, \quad \implies \quad \delta_2(2, \dots, 2) = -V_0(2, \dots, 2) = \delta_0^2(1, \dots, 1),$$

$$V_2(4, \dots, 4) = 0, \quad \implies \quad \delta_3(2, \dots, 2) = \delta_2(2, \dots, 2) = \delta_0^2(1, \dots, 1).$$

From these two conditions and (5.15) we see that (5.14) is valid.

One can use the vanishing conditions of $V_j(4, \dots, 4, 1, \dots, 1)$ or $V_j(4, \dots, 4, 3, \dots, 3)$ to write down a second-order difference equation for $\delta_j(2, \dots, 2, 1, \dots, 1)$, for $3 \leq j \leq r-3$.

The two initial conditions are:

$$\begin{aligned}\delta_0(2, \dots, 2, 1, \dots, 1) &= 0, \quad \implies \quad \delta_2(2, \dots, 2, 1, \dots, 1) = -V_0(2, \dots, 2, 1, \dots, 1), \\ V_2(4, \dots, 4, 1, \dots, 1) &= 0.\end{aligned}$$

Factoring out $\delta_j(2, \dots, 2, 0, \dots, 0) \prod_{a \leq i < b} \delta_{ab,0}(1, 1)$ from the equations and the conditions, and comparing them to $V_j(0, \dots, 0, 1, \dots, 1)$, we see that (5.13) is valid.

One can write down the second-order linear difference equations for $\delta_j(1, \dots, 1)$, for $3 \leq j \leq r - 3$. They are in the same form as in the $A_r^{(1)}$ case. Choosing the y 's in (3.9) to be $y_a^{\sigma_a} = e^{-i\sigma_a \vartheta_a}$ and comparing to (5.12), one sees that (5.12) satisfies the difference equation. However, this does not prove (5.12) uniquely; one still has to verify that the boundary conditions are indeed satisfied.

6 Concluding Remarks

Hirota's method has been used to construct exact solutions for affine Toda theories. We have endeavored systematically to explore this iterative method and although we do not claim completeness, we have presented new, explicit multi-soliton solutions for all the cases discussed, including the detailed procedures for how they were obtained. The double-soliton solutions given can be proved explicitly, while general N -soliton solutions are hard to verify directly. Scattering solutions of E_6 , E_7 or E_8 affine Toda solitons can also be obtained in this way, due to the lengthiness of the expressions, they are not included here. In addition to the simply-laced loop algebras which we discussed in detail, we have also given some examples of multi-soliton solutions of non-simply-laced ones. These can be solved directly using the iterative procedures, or indirectly, as we have done, by twisting the simply-laced solutions.

For simply-laced affine TFTs, the $\kappa = 2$ or 3 solitons are actually composites of more fundamental $\kappa = 1$ solitons, and can be obtained by taking the zero-velocity limit of some

scattering solutions, even though they have single-soliton form. For non-simply-laced affine TFTs, the $\kappa = 2$ or 3 solitons are just as fundamental as $\kappa = 1$ solitons.

After this phase of our work was completed, additional recent work by us [19] cast considerable doubt that many of the soliton solutions actually have real energy density, contrary to the hopes expressed by others [1, 4]. While not negating the interest of the multi-soliton solutions presented here vis-à-vis the knowledge they give us about the detailed classical spectrum of the ATFTs per se, nevertheless, these problems about real energy densities may cause difficulties in considering the physical applications of some of the theories.

The single-soliton solutions have a continuous parameter ξ , which is related to the shape and topological charge of the soliton. For most choices of ξ -value, the energy density is a complex function of x and t . By demanding the energy density of a single-soliton be real, then ξ can only take discrete values. However, the reality is not always preserved in the scattering. All this will be explored further in a future publication [19].

After our work was completed, we received the paper of Olive *et. al* [10], in which a different method, based on Vertex operators, for finding soliton solutions for TFT's is developed. It will be interesting to compare explicit results found with their method to our results.

References

- [1] T. Hollowood, *Nucl. Phys.* **B384** (1992) 523.
- [2] T.J. Hollowood, *Int. J. Mod. Phys.* **A8** (1993) 947.
- [3] Timothy J. Hollowood and Paul Mansfield, *Nucl. Phys.* **B226** (1989) 73.
- [4] Jonathson M. Evans *Nucl. Phys.* **B390** (1993) 225.

- [5] O. Babelon and L. Bonora, *Phys. Lett.* **244B** (1990) 220.
- [6] N. J. MacKay and W. A. McGhee, *Int. J. Mod. Phys.* **A8** (1993), 2791.
- [7] H. Aratyn, C. P. Constantinidis, L. A. Ferreira, J. F. Gomes, A. H. Zimerman, “Hirota’s solitons in the Affine and Conformal Affine Toda Models”, IFT-P/052/92, UICHEP-TH/92-18, hep-th/9212086.
- [8] H. C. Liao, D. I. Olive and N. Turok, *Phys. Lett* **B298** (1993) 95.
- [9] D. I. Olive, N. Turok and J. W. R. Underwood, “Solitons and the Energy-Momentum tensor for Affine Toda field theory”, Imperial/TP/91-92/35, SWAT/3.
- [10] D. I. Olive, N. Turok and J. W. R. Underwood, “Affine Toda Solitons and Vertex Operators”, IC/92-93/29, PUP-PH-93/1392, SWAT/92-93/5.
- [11] M. A. C. Kneipp and D. I. Olive, “Crossing and Antisolitons in Affine Toda Theories”, SWAT/92-93/6.
- [12] J. W. R. Underwood, “On the Topological Charges of Affine Toda Solitons”, Imperial/TP/92-93/38.
- [13] A. B. Zamolodchikov, *Int. J. Mod. Phys.* **A3** (1988) 743; *Int. J. Mod. Phys.* **A4** (1989) 4235; V. A. Fateev and A. B. Zamolodchikov, *Int. J. Mod. Phys.* **A5** (1990) 1025.
- [14] R. Hirota, “Direct methods in soliton theory”, in *Soliton* page 157; R.K. Bullough and P.S. Caudrey eds., (Springer–Verlag, Berlin 1980).
- [15] P. Christe and G. Mussardo, *Int. J. Mod. Phys.* **24** (1990) 4581. H.W. Braden, E. Corrigan, P.E. Dorey and R. Sasaki, *Nuclear Physics* **B356** (1991) 469; *ibid* **B338** (1990) 689.

- [16] D.I. Olive and N. Turok, *Nuclear Physics* **B215** [FS7] (1983) 470.
- [17] T.J. Hollowood, “Quantum soliton mass corrections in $SL(N)$ affine Toda field theory”,
OUTP-92-19P, hep-th 9209024.
- [18] William A. McGhee, “The Topological Charges of the $a_n^{(1)}$ Affine Toda Solitons”, DTP-
93-35, hep-th/9307035.
- [19] Z. Zhu and D. G. Caldi, “Reality of Complex Affine Toda Solitons”, in preparation.